

THE INTEGRAL CORE OBSERVING PROGRAMME

C. Winkler ¹, N. Gehrels ², N. Lund ³, V. Schönfelder ⁴ and P. Ubertini ⁵

(1) *ESA/ESTEC, Noordwijk, The Netherlands*, (2) *NASA/GSFC, Greenbelt, USA*, (3) *DSRI, Copenhagen, Denmark*, (4) *MPE, Garching, Germany*, (5) *IAS, Rome, Italy*

ABSTRACT The Core Programme of the INTEGRAL mission is defined as the portion of the scientific programme covering the guaranteed time observations for the INTEGRAL Science Working Team. This paper describes the current status of the Core Programme preparations and summarizes the key elements of the observing programme.

KEYWORDS: INTEGRAL; nucleosynthesis; compact objects; high energy transients.

1. INTRODUCTION

Scientific observing time for the observing programme during the nominal (2 yr) and extended (3 yr) mission phases of INTEGRAL is divided into the open time for the General Observer (General Programme; year 1: 65%, year 2: 70%, year 3+: 75%) and the guaranteed time for the INTEGRAL Science Working Team (ISWT) (Core Programme; year 1: 35%, year 2: 30%, year 3+: 25%). Observing time for the General Programme will be allocated to the scientific community during the AO process. The Core Programme will consist of three elements (see also Gehrels et al. 1997, Ubertini et al. 1997a): (i) Frequent scans of the Galactic plane (Galactic Plane Survey, GPS) (ii) Deep exposure of the Galactic central radian (Galactic Central Radian Deep Exposure, GCDE) (iii) Pointed observations of selected sources. The data from the Core Programme belong to the ISWT for the usual proprietary period of one year, after which the data will become public.

2. SCIENCE OBJECTIVES OF THE CORE PROGRAMME

The scanning of the Galactic plane (GPS) will be mainly done for two reasons: the most important one is to provide frequent monitoring of the plane in order to detect transient sources because the gamma-ray sky in the INTEGRAL energy range is dominated by the extreme variability of many sources. The scans would find sources in high state (outburst) which warrant possible scientifically important follow-up observations (Target of Opportunity [TOO] observation). The second reason is to build up time resolved maps of the Galactic plane in continuum and diffuse line emission such as ²⁶Al and 511 keV with modest exposure. Compton GRO and SIGMA have been detecting Galactic compact sources of several different categories/groups which include X-ray binaries (e.g. X-ray novae, Be binary pulsars) and in particular superluminal sources (GRS 1915+105 and GRO J1655–40). The occurrence rates for events that INTEGRAL can observe is about 2 events/year for

each of these classes, where pointing constraints due to the fixed solar arrays have been taken into account. The important time scales for the transient outbursts vary significantly from class to class and from event to event, but a typical duration of an event is 1 - 2 weeks and a typical variability time scale is of the order of 1 day. The scans will be performed once a week by performing a “slew - and stare” manoeuvre of the spacecraft along the visible (accessible) part of the Galactic plane with latitude extent $\pm 10^\circ$. The accessible part of the Galactic plane depends on viewing constraints including the solar aspect angle (40° during first two years, 30° thereafter) due to the fixed solar arrays, and on the season of the year. The angular distance between subsequent exposures (965 s each) along the scan path is 6° . The scan will be performed as a sawtooth with inclination of 21° with respect to the Galactic plane, each subsequent scan is shifted by 27.5° in galactic longitude.

The deep exposure of the central Galactic radian (GCDE) is driven by the following objectives: mapping line emission from nucleosynthesis radioisotopes (e.g. ^{26}Al , ^{44}Ti , 511 keV), mapping continuum emission of the Galactic ridge, and performing deep imaging and spectroscopic studies of the central region of the Galaxy. Several interesting emission regions in or near the Galactic plane have been identified using CGRO OSSE and COMPTEL: these include the ^{26}Al (7×10^5 years half life) mapping the sites of nucleosynthesis in the past million years in the Galaxy (Diehl et al. 1995), and ^{44}Ti (half life ~ 60 year) which has been detected by COMPTEL from the Cas A SNR (Iyudin et al. 1994). OSSE mapping of the positron - electron annihilation radiation at 511 keV shows a central bulge, emission in the Galactic plane and an enhancement of extension of emission at positive latitudes above the Galactic centre (Purcell et al. 1997). Other isotopes such as ^{60}Fe produce lines which could be detected by INTEGRAL. The origin of the clumpy structure of the COMPTEL observed ^{26}Al maps and the ^{44}Ti emission from hidden supernovae are key targets of INTEGRAL research. The INTEGRAL deep exposure will also study the continuum gamma-ray and hard X-ray emission from the Galactic plane. This “galactic ridge” is concentrated in a narrow band with a latitude extent of $\sim 5^\circ$ and a longitude extent of $\pm 40^\circ$ (Gehrels & Tueller 1993, Valinia & Marshall 1998, Strong et al. 1999). The exact distribution and spectrum of the ridge emission is not well known. The origin is thought to be Bremsstrahlung from cosmic ray electrons, but this is also not fully established. INTEGRAL will be able to map the emission with high sensitivity and high angular resolution. This should allow the removal of the point-source origin of the emission so that the spectrum can be determined with high confidence. The GCDE will resolve isolated point sources with arcmin location and provide source spectra with high energy resolution. At least 90 sources known as X- and gamma-ray emitters are contained in the region at ~ 10 keV and GRANAT/SIGMA and other earlier experiments have found this region to be filled with a number of highly variable and transient sources (Vargas et al. 1997, Ubertini et al. 1997). Many are thought to be compact objects in binary systems undergoing dynamic accretion. However, even for some of the brightest sources in the region (1E 1740.7–2942, GRS 1758–258) the detailed nature of the systems are not known. INTEGRAL will study the faintest sources with high angu-

lar resolution allowing multi-source monitoring within its wide field of view during single pointings. Also interesting will be searches for gamma-ray emission from SgrA*, at the Centre of the Galaxy (Sunyaev et al. 1993). The central radian of the Galaxy will be observed using a rectangular pointing grid of $11 \times 31 = 341$ pointings with a pitch of 2° . The grid covers the celestial region between $-30^\circ \leq l \leq +30^\circ$ and $-10^\circ \leq b \leq +10^\circ$. With 3 minutes slew duration for 2° and 1000 s exposure per point, about 4×10^5 s will be required to scan the grid once. Within the total time allocation for the GCDE, the grid scan will be performed 12 times per year. Remaining Core Programme observing time not spent on GPS and GCDE will be devoted to dedicated pointings on individual sources. These sources include galactic black hole candidates, neutron stars, AGN and some time will also be set aside to be able to perform scientifically important follow-up observations on TOO's which have been detected during GPS and GCDE or where the trigger has been observed by optical telescopes (SN and Novae). The greatest advantage of INTEGRAL, as compared to other high energy missions, is its high spectral resolution and high sensitivity at energies above 100 keV. The strongest point for INTEGRAL to response to a TOO call is the potential to detect transient high energy emission features (e.g. 511 keV, its scattering feature at 170 keV; 480 keV (${}^7\text{Li}$); MeV bump; flares from Blazars). The high energy cutoff or spectral break as a function of time may provide critical information to our understanding of the emission mechanism and system parameters. Concerning the high energy continuum, several pointings for each flare or outburst are required to achieve meaningful science return since the spectral evolution instead of simply the spectral shape is the main objective here. Separation between the exposures depends on the time scale of the event. The final list of candidate sources will be published, together with all details of the other Core Programme elements, in the first AO for INTEGRAL observations, which is scheduled for release at the end of 1999. General Observers cannot propose for targets included in the Core Programme list of candidate sources, unless the scientific objectives of their proposals are of a clearly different nature.

The orbit characteristics for INTEGRAL on PROTON (baseline) and ARIANE 5 (backup) are summarised in Winkler (1999) and Carli (1999). The allocation of observing time has been driven by highest scientific priority which has been given to GPS and GCDE and is presently scheduled for the first year of operations (baseline orbit) as follows: GCDE: 4.8×10^6 s; GPS: 2.3×10^6 s; Pointed observations: 2.2×10^6 s assuming an observation efficiency of 85%. It is planned to keep the allocations constant for GCDE and GPS during the year 2 of operations (and for GCDE during year 3 to 5).

3. EXPOSURE TIMES AND SENSITIVITIES

Due to the large coded field-of-views for SPI (FCFOV 16° across) and IBIS (FCFOV $9^\circ \times 9^\circ$), a single "staring" point on a GPS scan will not only receive the nominal exposure per pointing (965 s): While this exposure point is still located within the partially coded field-of-view (PCFOV, coverage to zero response), addi-

tional exposure will be collected as the spacecraft (i.e. the co-aligned instruments) is pointing at close neighbouring positions of that single exposure point.

For SPI and IBIS, a single “staring” point will therefore receive ~ 3000 s of exposure per scan, while JEM-X and OMC – because of the smaller FOV’s – will accumulate 965 s per point/scan. The 3σ sensitivity which can be obtained for one “staring” point during a single scan is therefore 10 mCrab @ 100 keV (IBIS), 750 mCrab @ 1 MeV (SPI) and 6 mCrab @ 10 keV (JEM-X). This will allow detection of transient sources throughout the Galaxy. During the first two years the total exposure per point per year is $\sim 9 \times 10^4$ s for SPI and IBIS. For JEM-X and OMC the exposure per point and year amounts to 2.7×10^4 s. The area covered by the GCDE will receive a net exposure of 4×10^6 s per year which will give SPI a 3σ sensitivity of $\sim 5 \times 10^{-6}$ ph/(cm²s) for narrow lines in the 100 keV - 2 MeV region sufficient for mapping and for detailed line shape studies at 1809 keV (Gehrels & Chen 1996) of the bright ²⁶Al “hot spots” ($\sim 3 \times 10^{-5}$ ph/(cm² s)) as detected by COMPTEL. The continuum sensitivities (3σ) in the 100 keV to 1 MeV range would be 5×10^{-7} to 4×10^{-8} ph/(cm² s keV) (SPI) and $(2 \text{ to } 1) \times 10^{-7}$ ph/(cm² s keV) (IBIS) while JEM-X would achieve 10^{-5} to 10^{-6} ph/(cm² s keV) in the 3 to 30 keV range. Pointed observations, e.g. nucleosynthesis studies of Cygnus and Vela regions require typically $\sim 10^6$ s while Galactic compact (transient) objects could be observed with 10^5 s or less per repeated observation. Given the high sensitivity of INTEGRAL during the GPS (and GCDE), follow-up observations for Galactic TOO’s - like the superluminal jet source GRS 1915+105 or Cyg X–1 - could be triggered if the 100 keV flux would exceed a few hundred mCrab, easily detectable in one GPS exposure. Supernovae and Novae observations would be possibly done repeatedly at various times after onset, to study (e.g. for SNII) early observations of radioactivity from Co breaking through the SN envelope and late observations of ⁴⁴Ti.

ACKNOWLEDGMENTS We thank G. Skinner and P. Connell for simulation studies.

REFERENCES

- Carli R et al., 1999, these proceedings
- Diehl R., et al., 1995, A&A 298, 445
- Gehrels N., Tueller J., 1993, ApJ 407, 597
- Gehrels N., Chen W., 1996, A&AS 120, 331
- Gehrels N., et al., 1997, ESA SP-382, 587
- Iyudin A., et al., 1994, A&A 284, L1
- Purcell W., et al., 1997, ApJ 491, 725
- Strong A., et al., 1999, these proceedings
- Sunyaev et al., 1993, ApJ 407, 606.
- Ubertini P., et al., 1997, Proc. 4th CGRO Symposium, AIP 410, 1527
- Ubertini P., et al., 1997a, Proc. Workshop on ‘ASM and GRB emission’
- Valinia A., Marshall F., 1998, ApJ 505, 134
- Vargas M., et al., 1997, ESA SP-382, 129
- Winkler C., 1999, these proceedings